


$$t/\tau_{td} = 2.3$$

Formation of sunspots

[turbulence] [stratification] [magnetic field]

Illa R. Losada

Nordita

Stockholm University

1st SOLARNET Spring School: "Introduction to Solar Physics"
March 2014



Flux tubes

Formation of sunspots

Illa R. Losada

Flux tubes

Back to equations

MHD
DNS and MFS
DNS
MFS
MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results
Horizontal field
Vertical field
Parameters study

Corona: two layers model

Conclusions




Figure 1 : Caligari et al. (1995)

Flux tubes: problems.

Flux tubes

Back to equations

MHD
DNS and MFS
DNS
MFS
MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results
Horizontal field
Vertical field
Parameters study

Corona: two layers model

Conclusions

- ▶ Magnetic field needed at the bottom $\approx 10^5$ Gauss
 - ▶ $B_{\text{eq}} = 10^4$ Gauss; $B_{\text{eq}} = \sqrt{\mu_0 \rho} u_{\text{rms}}$
 - ▶ buoyancy dominates
 - ▶ “tild angle” (between connecting line of 2 polarities and equator)
 - ▶ emergence latitude

Flux tubes: problems.

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study


Corona: two layers model

Conclusions

- ▶ Magnetic field needed at the bottom $\approx 10^5$ Gauss
- ▶ Importance of the tachocline?
Also cycles in totally convective stars.

Flux tubes: problems.

- ▶ Magnetic field needed at the bottom $\approx 10^5$ Gauss
- ▶ Importance of the tachocline?
- ▶ Helioseismology (wave-speed anomalies)



Flux tubes

Back to equations

MHD
DNS and MFS
DNS
MFS
MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results
Horizontal field
Vertical field
Parameters study

Corona: two layers model

Conclusions

Flux tubes: problems.

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers model

Conclusions

- ▶ Magnetic field needed at the bottom $\approx 10^5$ Gauss
- ▶ Importance of the tachocline?
- ▶ Helioseismology
- ▶ Avoiding distortion while raising in a turbulent media.
 - ▶ expansion of the tubes along their raise
 - ▶ thin tubes: dynamics of the surrounding medium dominates

Solve MHD equations

Let's go back to the equations!

Formation of
sunspots

Illa R. Losada

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

Solve MHD equations

Formation of
sunspots

Illa R. Losada

Let's go back to the equations!

MHD equations

- ▶ Magnetohydrodynamics (MHD): equations for the dynamics of the plasma.

- ▶ Goal of simulations: solve MHD equations.

- ▶ Not possible to use solar parameters:

$$\text{Re} \approx \frac{UL}{\nu} \approx 10^{10} - 10^{15}$$

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

Solve MHD equations

Formation of
sunspots

Illa R. Losada

Let's go back to the equations!

MHD equations

- ▶ Magnetohydrodynamics (MHD): equations for the dynamics of the plasma.
- ▶ Goal of simulations: solve MHD equations.
- ▶ Not possible to use solar parameters:
$$\text{Re} \approx \frac{UL}{\nu} \approx 10^{10} - 10^{15}$$

Simulations

- ▶ Direct Numerical Simulations (DNS)
- ▶ Mean-field Simulations (MFS)
- ▶ Large Eddy Simulations (LES)

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

MFS and DNS: How to cook paella!

Formation of
sunspots

Illa R. Losada

Mean Field Simulations (MFS)

Direct Numerical Simulations (DNS)

- ▶ Quantities = averaged + fluctuations: $F = \overline{F} + f$
- ▶ Approximations: add or subtract terms in the equations.
- ▶ Control the physics.

- ▶ Solve full equations.
- ▶ Approximations: only in resolution.
- ▶ No control.



Simulations done with Pencil Code
(<http://pencil-code.googlecode.com>)

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

Full MHD equations.

Direct Numerical simulations (DNS):

- ▶ Continuity equation: $\frac{\partial \rho}{\partial t} = -\nabla \cdot (\rho \mathbf{U})$
- ▶ Momentum equation:
$$\frac{D\mathbf{U}}{Dt} = -\nabla p + \mathbf{J} \times \mathbf{B} + \rho \mathbf{g} + \rho \nu (\nabla^2 \mathbf{U} + \frac{1}{3} \nabla \nabla \cdot \mathbf{U} + 2 \mathbf{S} \cdot \nabla \ln \rho)$$
- ▶ Induction equation: $\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{U} \times \mathbf{B} - \eta \mu_0 \mathbf{J})$

Formation of
sunspots

IIIa R. Losada

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

Full MHD equations.

Direct Numerical simulations (DNS):

- ▶ Continuity equation: $\frac{\partial \rho}{\partial t} = -\nabla \cdot (\rho \mathbf{U})$
- ▶ Momentum equation:
$$\frac{D\mathbf{U}}{Dt} = -\nabla p + \mathbf{J} \times \mathbf{B} + \rho \mathbf{g} + \rho \nu (\nabla^2 \mathbf{U} + \frac{1}{3} \nabla \nabla \cdot \mathbf{U} + 2 \mathbf{S} \cdot \nabla \ln \rho)$$
- ▶ Induction equation: $\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{U} \times \mathbf{B} - \eta \mu_0 \mathbf{J})$

Setup:

- ▶ Forcing, $k_f/k_1 = 15$
(control scale separation)
- ▶ Strong stratification:
density contrast ≈ 535
- ▶ $B_0/B_{\text{eq}0} = 0.05$ (in range)
- ▶ $64^3 \times 128$ mesh-points
- ▶ $\text{Re}_M \approx \frac{UL}{\eta} = 6$

$$\Delta \bar{\mathbf{B}}/B_{\text{eq}0} \text{ (Brandenburg et al. 2011)}$$

Mean-Field MHD equations.

Quantities = averaged + fluctuations: $F = \overline{F} + f$

Mean field simulations (MFS):

► Continuity equation: $\frac{D\overline{\rho}}{Dt} = -\overline{\rho} \nabla \cdot \overline{\mathbf{U}}$

► Momentum equation:

$$\frac{D\overline{\mathbf{U}}}{Dt} = -c_s^2 \nabla \ln \overline{\rho} + \mathbf{g} + \overline{\mathcal{F}}_M + \overline{\mathcal{F}}_K$$

► Induction equation:

$$\frac{\partial \overline{\mathbf{B}}}{\partial t} = \nabla \times (\overline{\mathbf{U}} \times \overline{\mathbf{B}} + \overline{\mathbf{u} \times \mathbf{b}}) + \eta \nabla^2 \overline{\mathbf{B}}$$

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

From induction equation.

$$\frac{\partial \overline{\mathbf{B}}}{\partial t} = \nabla \times (\overline{\mathbf{U}} \times \overline{\mathbf{B}} + \overline{\mathbf{u} \times \mathbf{b}}) + \eta \nabla^2 \overline{\mathbf{B}}$$

α -effect (Steenbeck et al. 1966; Moffat 1978; Krause & Rädler 1980)

- ▶ Isotropic case: $\overline{\mathbf{u} \times \mathbf{b}} = \alpha \overline{\mathbf{B}} - \eta_t \overline{\mathbf{J}}$
- ▶ If $\alpha \neq 0 \rightarrow$ generate a $\overline{\mathbf{B}}$
- ▶ η_t : turbulent diffusivity.

Responsible for the Sun's large-scale field.

From momentum equation.

$$\frac{D\bar{\mathbf{U}}}{Dt} = -c_s^2 \nabla \ln \bar{\rho} + \mathbf{g} + \bar{\mathcal{F}}_M + \bar{\mathcal{F}}_K$$

Mean Lorentz Force, $\bar{\mathcal{F}}_M$:

$$\bar{\rho} \bar{\mathcal{F}}_M = -\nabla(\overline{\rho \mathbf{u} \mathbf{u}} + \dots) = \bar{\mathbf{J}} \times \bar{\mathbf{B}} + \frac{1}{2\mu_0} \nabla(q_p \bar{\mathbf{B}}^2) + \dots$$

$\bar{\mathbf{J}} = \nabla \times \bar{\mathbf{B}} / \mu_0 = -\frac{1}{2} \nabla \bar{\mathbf{B}}^2 + \bar{\mathbf{B}} \cdot \nabla \bar{\mathbf{B}}$: mean current density,
 $\frac{1}{2} \nabla(q_p \bar{\mathbf{B}}^2)$: turbulent contribution


$$\bar{\rho} \bar{\mathcal{F}}_M = -\frac{1}{2} \nabla[(1 - q_p) \bar{\mathbf{B}}^2] + \bar{\mathbf{B}} \cdot \nabla[(1 - q_s) \bar{\mathbf{B}}]$$

Pressure. NEMPI.

Effective magnetic pressure (Kemel et. al 2012):

(effects of turbulence on the mean Lorentz force)

$$P_{eff} = \frac{1}{2}(1 - q_p)\overline{\mathbf{B}}^2 / \overline{B}_{eq}^2$$



Negative Effective Magnetic Pressure Instability (NEMPI)

- ▶ Regions below the minimum value of $P_{eff} \rightarrow$ NEMP
- ▶ NEMP + strong stratification ($|\nabla \ln \rho| > |\nabla \overline{\mathbf{B}}|$) \rightarrow NEMPI¹
- ▶ Magnetic field suppress turbulence \rightarrow structures sink!

¹ Predicted: Kleeorin et al. 1989, 1990; Kleeorin & Rogachevskii 1994; Kleeorin et al. 1996; Rogachevskii & Kleeorin 2007. Confirmed: Brandenburg et al. 2011

Normalized effective magnetic pressure:

$$\mathcal{P}_{\text{eff}} = \frac{1}{2}(1 - q_p)\beta^2, \quad (1)$$

$q_p(\beta)$ approximated by (Kemel et al. 2012a):

$$q_p(\beta) = \frac{q_{p0}}{1 + \beta^2/\beta_p^2} = \frac{\beta_\star^2}{\beta_p^2 + \beta^2}, \quad (2)$$

q_{p0} , β_p , and $\beta_\star = \beta_p q_{p0}^{1/2}$: constants.

\mathcal{P}_{eff} has a minimum value \mathcal{P}_{min} at β_{min} , related with the parameters:

$$\beta_p = \beta_{\text{min}} / \sqrt{-2\mathcal{P}_{\text{min}}}, \quad \beta_\star = \beta_p + \sqrt{-2\mathcal{P}_{\text{min}}}. \quad (3)$$

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

Velocity field and NEMPI

Forced Velocity field

NEMPI.

$$P = P_{gas} + P_{eff}$$

Formation of
sunspots

Illa R. Losada

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

MFS vs. DNS.

Formation of sunspots

Ilia R. Losada




Figure 2 : Time evolution of the magnetic field and effective magnetic pressure in mean-field simulations (Kemel et al. 2012).




Figure 3 : Time evolution of the mean (y-averaged) magnetic field and effective magnetic pressure in direct numerical simulations (Kemel et al. 2012).

High degree of predictive power of MFS.

Different scales because of different input parameters

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers model

Conclusions

Previous results.

- ▶ NEMP and NEMPI described in DNS and MFS ².
- ▶ NEMPI observed in DNS and MFS ³
- ▶ Parameters values studied for maximize the growth strength and time:
 - ▶ $k_f/k_1 = 30$ (scale separation ratio)
 - ▶ $Re \equiv u_{rms}/\nu k_f = 36$
 - ▶ $P_m = \nu/\eta = 0.5$.
 - ▶ $Re_M = P_m Re = 18$
- ▶ Effects of rotation.

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

My contribution:

- ▶ Effects of rotation
- ▶ Effects of stratification
- ▶ Effects of corona: 2 layers model

²Kleeorin et al. 1989, 1989, 1996; Kleeorin & Rogachevskii 1994;

Rotation. Structures evolution

(Rotational effects on the negative magnetic pressure instability (2012, A&A,548,A49))

(Competition of rotation and stratification in flux concentrations (2013, A&A, 556, A83))

$$Co = 2\Omega\tau_c = 2\Omega/u_{\text{rms}}k_f$$

$$Co = 0.006$$

$$Co = 0.03$$

$$Co = 0.06$$

Simulations done with Pencil Code (<http://pencil-code.googlecode.com>)

Formation of
sunspots

Illa R. Losada

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

MFS and DNS results.

Formation of sunspots

Ilia R. Losada




Figure 4 : MFS (i): $q_{p0} = 20$ and $\beta_p = 0.167$; MFS (ii): $q_{p0} = 32$ and $\beta_p = 0.058$.

$$B_0/B_{eq0} = 0.05. \quad q_p = \frac{\beta_{*}^2}{(\beta_p^2 + \beta_{*}^2)} \quad \beta = \frac{\bar{B}}{B_{eq0}} \quad \lambda_{*0} \equiv \beta_{*} u_{rms}/H_p$$

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers model

Conclusions

Polytropic stratification. Horizontal imposed field (MFS).

Formation of
sunspots

Ilia R. Losada

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions




Figure 5 : \overline{B}_y in the kinematic growth phase for different γ and β_0 in the presence of a horizontal field using the perfect conductor boundary condition.

Polytropic stratification. Vertical imposed field (MFS). Evolution.

$$\gamma = 1$$

$$\gamma = 5/3$$

$$\beta_0 = 0.02$$

Formation of sunspots

Illa R. Losada

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers model

Conclusions

Vertical field. Sunspot-like structure

Formation of sunspots

Ilia R. Losada




Figure 6 : Cuts of $B_z/B_{eq}(z)$ in the xy plane at the top boundary ($z/H_{\rho 0} = 1.2$) and the xz plane through the middle of the spot at $y = 0$ for $\gamma = 5/3$ and $\beta_0 = 0.05$. In the xz cut, we also show magnetic field lines and flow vectors obtained by numerically averaging in azimuth around the spot axis.

Flux tubes

Back to equations

MHD
DNS and MFS
DNS
MFS
MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers model

Conclusions

Vertical imposed field (DNS vs MFS). Evolution.

DNS

MFS

$$\beta_0 = 0.05$$

Formation of
sunspots

Illa R. Losada

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

Parameters dependence study in polytropic stratification

Formation of sunspots

Illa R. Losada

Flux tubes

Back to equations

MHD
DNS and MFS
DNS
MFS
MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results
Horizontal field
Vertical field
Parameters study

Corona: two layers model

Conclusions




Figure 7 : Effective magnetic pressure obtained from DNS in a polytropic layer with different γ for horizontal (H, red curves) and vertical (V, blue curves) mean magnetic fields.




Figure 8 : Parameters q_{p0} , β_p , and β_* for the function $q_p(\beta)$ versus γ for horizontal (red line) and vertical (blue line) mean magnetic fields.

Corona: two layers model

Bipolar magnetic structures driven by stratified turbulence with a coronal envelope (2013, ApJ Lett 777, L37; arXiv:1308.1080)

Bipolar magnetic structures.

$$256^2 \times 512, kf/k1 = 30, Re_M = 18$$

Formation of
sunspots

Illa R. Losada

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field









Vertical field

Parameters study

**Corona: two layers
model**

Conclusions

Bipolar magnetic structures



Formation of sunspots

Ilia R. Losada

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers model

Conclusions

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

Conclusions

Parker instability vs. NEMPI

	NEMPI	Parker Instability
Turbulence	turbulent	non turbulent
Scale	sufficient many eddies	small
Stratification	continuous strat. B	non uniform & initially separated
Energy source	turbulent energy	gravitational field
Initial field	smooth: $H_B^{-1} \equiv \nabla \ln B $ small	structured: $ \nabla \ln B $ large
Density variation	$H_\rho^{-1} \equiv \nabla \ln \rho $ large	$H_\rho^{-1} \equiv \nabla \ln \rho $ small
Instability criteria	$\frac{H_B}{H_\rho} \gg 1$	$\frac{H_B}{H_\rho} \ll 1$

Conclusions.

Rotation.

NEMPI depends on the ratio between rotation and turbulence.

- ▶ Turnover time: $\tau \approx 2\text{hours}$
- ▶ On the Sun: only upper-most layers (super-granulation layer $\tau \sim 1\text{day}$)

NEMPI + Dynamo:

- ▶ Coupled system for $Co > 0.1$
- ▶ Direct comparison with NEMPI growth rate not possible.

Conclusions.

Polytropic EoS:

- ▶ NEMPI develops in the uppermost layers.
- ▶ The parametrization depends on the polytropic index.
- ▶ Isothermal models applicable locally.
- ▶ No “potato-sack” effect with vertical fields.

Coronal envelope:

- ▶ Emergence and decaying of bipolar magnetic region.
- ▶ The 2 polarities tend to separate.

Solar activity as a surface phenomenon.

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers
model

Conclusions

What is next?.

- ▶ Effects of rotation in a vertical field configuration.
- ▶ NEMPI dependence on Re .
- ▶ MFS studies of the two-layer model.
- ▶ Polytropic atmosphere in the two-layer model.
- ▶

Illa R. Losada

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers model

Conclusions

Sunspot formation: theory, simulations and observations



Workshop
9–13 March 2015

Coordinators: Illa R. Losada,
Göran Scharmer, Nishant Singh,
Axel Brandenburg

www.nordita.org/sunspots2015

S.O.C.: Axel Brandenburg (PI, Nordita), Mark Cheung (Lockheed, Palo Alto), Laurent Gizon (MPS, Göttingen), Petri Käpylä (Helsinki University), Alexander Kosovichev (Big Bear), Åke Nordlund (NBI), Matthias Rempel (HAO/NCAR), Göran Scharmer (Co-PI, Stockholm University), Alexandra Tritschler (NSO, New Mexico), and Jörn Warnecke (MPS, Göttingen).

L.O.C.: Illa R. Losada (PI, Nordita and Stockholm University), Sarah Jabbari (Nordita and Stockholm University), Bidya Binay Karak (Nordita), Hiva Pazira (Stockholm University), Nishant Singh (Nordita), Elizabeth Yang (Nordita, conference secretary).

Flux tubes

Back to equations

MHD

DNS and MFS

DNS

MFS

MFS vs DNS

First results

Rotation

Results

Polytropic EoS

Results

Horizontal field

Vertical field

Parameters study

Corona: two layers model

Conclusions

$$\frac{\partial g_{ik}}{\partial x^m} = -g_{ik} \Gamma_{mm}^m$$

$$\frac{\partial}{\partial x^p} (\delta_j^i) = 0 \Rightarrow \frac{\partial}{\partial x^p} (g^{ik} g_{kj}) = 0$$

$$\Rightarrow g^{ik} \frac{\partial}{\partial x^p} g_{kj} + \left(\frac{\partial}{\partial x^p} g^{ik} \right) g_{kj} = 0$$

$$g^{ji} g_{kj} \frac{\partial g^{ik}}{\partial x^p} = -g^{ji} g^{ik} \frac{\partial g_{kj}}{\partial x^p}$$

$$\delta_k^j \frac{\partial g^{ik}}{\partial x^p} = -g^{ji} g^{ik} [k p, j] - g^{ji} g^{ik} [j p, k]$$

$$\frac{\partial g^{ik}}{\partial x^p} = -g^{ik} \Gamma_{kp}^j - g^{ji} \Gamma_{jp}^i \quad \square$$



Thank you for your time.